

LAMPIRAN 8
UJI NORMALITAS

Case Processing Summary

| | Valid | | Cases Missing | | Total | |
|---------------|-------|---------|---------------|---------|-------|---------|
| | N | Percent | N | Percent | N | Percent |
| TOTAL.DK | 40 | 100,0% | 0 | 0,0% | 40 | 100,0% |
| TOTAL.CBURDEN | 40 | 100,0% | 0 | 0,0% | 40 | 100,0% |

Descriptives

| | | Statistic | Std. Error | |
|----------------------------------|----------------------------------|-------------|------------|---------|
| TOTAL.DK | Mean | 66,5000 | 1,06639 | |
| | 95% Confidence Interval for Mean | Lower Bound | 64,3430 | |
| | | Upper Bound | 68,6570 | |
| | 5% Trimmed Mean | 66,4722 | | |
| | Median | 66,0000 | | |
| | Variance | 45,487 | | |
| | Std. Deviation | 6,74442 | | |
| | Minimum | 46,00 | | |
| | Maximum | 82,00 | | |
| | Range | 36,00 | | |
| | Interquartile Range | 7,75 | | |
| | Skewness | -,037 | ,374 | |
| | Kurtosis | 1,593 | ,733 | |
| | TOTAL.CBURDEN | Mean | 24,0000 | 1,81730 |
| 95% Confidence Interval for Mean | | Lower Bound | 20,3242 | |
| | | Upper Bound | 27,6758 | |
| 5% Trimmed Mean | | 23,9444 | | |
| Median | | 23,0000 | | |
| Variance | | 132,103 | | |
| Std. Deviation | | 11,49359 | | |
| Minimum | | 4,00 | | |
| Maximum | | 45,00 | | |
| Range | | 41,00 | | |
| Interquartile Range | | 16,75 | | |
| Skewness | | -,043 | ,374 | |
| Kurtosis | | -,930 | ,733 | |

Tests of Normality

| | Kolmogorov-Smirnov ^a | | | Shapiro-Wilk | | |
|---------------|---------------------------------|----|-------------------|--------------|----|------|
| | Statistic | df | Sig. | Statistic | df | Sig. |
| TOTAL.DK | ,105 | 40 | ,200 [*] | ,957 | 40 | ,131 |
| TOTAL.CBURDEN | ,099 | 40 | ,200 [*] | ,967 | 40 | ,284 |

*. This is a lower bound of the true significance.

a. Lilliefors Significance Correction