

## Lampiran 6. Hasil Analisis Deskriptif

### Descriptive Statistics

	N	Minimum	Maximum	Mean	Std. Deviation
umur	36	1.00	2.00	1.4167	.50000
masakerja	36	1.00	2.00	1.3889	.49441
penggunaanmasker	36	1.00	2.00	1.4167	.50000
Riwayat penyakit	36	1.00	2.00	1.3333	.47809
Valid N (listwise)	36				

## Lampiran 7. Hasil Analisis Chi Square

### Crosstabs

#### Umur \* kelainan Crosstabulation

Count

		kelainan		Total
		Normal	Gangguan	
umur	< 30 tahun	18	3	21
	> 30 tahun	8	7	15
Total		26	10	36

### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	4.573 <sup>a</sup>	1	.032		
Continuity Correction <sup>b</sup>	3.102	1	.078		
Likelihood Ratio	4.588	1	.032		
Fisher's Exact Test				.058	.039
Linear-by-Linear Association	4.446	1	.035		
N of Valid Cases	36				

a. 1 cells (25.0%) have expected count less than 5. The minimum expected count is 4.17.

b. Computed only for a 2x2 table

**Masa Kerja \* kelainan Crosstabulation**

Count

		kelainan		Total
		Normal	Gangguan	
masakerja	Lama	18	4	22
	Baru	8	6	14
Total		26	10	36

**Chi-Square Tests**

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2- sided)	Exact Sig. (1- sided)
Pearson Chi-Square	2.597 <sup>a</sup>	1	.107		
Continuity Correction <sup>b</sup>	1.512	1	.219		
Likelihood Ratio	2.557	1	.110		
Fisher's Exact Test				.140	.110
Linear-by-Linear Association	2.524	1	.112		
N of Valid Cases	36				

a. 1 cells (25.0%) have expected count less than 5. The minimum expected count is 3.89.

b. Computed only for a 2x2 table

**Penggunaan Masker \* kelainan Crosstabulation**

Count

		kelainan		Total
		Normal	Gangguan	
Pemakaian Masker	ya	18	3	21
	tidak	8	7	15
Total		26	10	36

**Chi-Square Tests**

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	4.573 <sup>a</sup>	1	.032	.058	.039
Continuity Correction <sup>b</sup>	3.102	1	.078		
Likelihood Ratio	4.588	1	.032		
Fisher's Exact Test					
Linear-by-Linear Association	4.446	1	.035		
N of Valid Cases	36				

a. 1 cells (25.0%) have expected count less than 5. The minimum expected count is 4.17.

b. Computed only for a 2x2 table

**Riwayat Penyakit \* kelainan Crosstabulation**

Count

		kelainan		Total
		Normal	Gangguan	
Riwayat Penyakit	tidak pernah	21	3	24
		5	7	12
Total		26	10	36

### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	8.377 <sup>a</sup>	1	.004		
Continuity Correction <sup>b</sup>	6.248	1	.012		
Likelihood Ratio	8.155	1	.004		
Fisher's Exact Test				.007	.007
Linear-by-Linear Association	8.144	1	.004		
N of Valid Cases	36				

a. 1 cells (25.0%) have expected count less than 5. The minimum expected count is 3.33.

b. Computed only for a 2x2 table

## Lampiran 8. Hasil Analisis Regresi Berganda

### Regression

#### Model Summary<sup>b</sup>

Model	R	R Square	Adjusted R Square	Std. Error of the Estimate
1	.521 <sup>a</sup>	.271	.203	.40552

a. Predictors: (Constant), sesaknapas, penggunaanmasker, umur

b. Dependent Variable: kelainan

#### ANOVA<sup>a</sup>

Model		Sum of Squares	df	Mean Square	F	Sig.
1	Regression	1.960	3	.653	3.972	.016 <sup>b</sup>
	Residual	5.262	32	.164		
	Total	7.222	35			

a. Dependent Variable: kelainan

b. Predictors: (Constant), sesaknapas, penggunaanmasker, umur

**Coefficients<sup>a</sup>**

Model	Unstandardized Coefficients		Standardized Coefficients	t	Sig.	
	B	Std. Error	Beta			
	1	(Constant)	.582			.236
	umur	-.234	.262	-.258	-.893	.379
	penggunaanmasker	.191	.163	.211	1.171	.250
	riwayatpenyakit	.567	.262	.597	2.165	.038

a. Dependent Variable: kelainan

**Residuals Statistics<sup>a</sup>**

	Minimum	Maximum	Mean	Std. Deviation	N
Predicted Value	.8723	1.6312	1.2778	.23663	36
Residual	-.63121	.89362	.00000	.38776	36
Std. Predicted Value	-1.713	1.494	.000	1.000	36
Std. Residual	-1.557	2.204	.000	.956	36

a. Dependent Variable: kelainan