


LAMP IRAN



REGRESSION

```

/MISSING LISTWISE
/STATISTICS COEFF OUTS R ANOVA COLLIN TOL
/CRITERIA=PIN(.05) POUT(.10)
/NOORIGIN
/DEPENDENT ROA
/METHOD=ENTER BOPO NIM Inflasi
/SCATTERPLOT=(*SDRESID ,*ZPRED)
/RESIDUALS DURBIN NORM(ZRESID)
/SAVE RESID.
    
```

Regression

Variables Entered/Removed^d

Model	Variables Entered	Variables Removed	Method
1	Inflasi, NIM, BOPO ^a		Enter

a. All requested variables entered.

b. Dependent Variable: ROA

Model Summary^d

Model	R	R Square	Adjusted R Square	Std. Error of the Estimate	Durbin-Watson
1	.921 ^a	.848	.834	.20745	1.869

a. Predictors: (Constant), Inflasi, NIM, BOPO

b. Dependent Variable: ROA

ANOVA^d

Model		Sum of Squares	df	Mean Square	F	Sig.
1	Regression	8.145	3	2.715	63.088	.000 ^a
	Residual	1.463	34	.043		
	Total	9.608	37			

a. Predictors: (Constant), Inflasi, NIM, BOPO

b. Dependent Variable: ROA


Coefficients^a

Model		Unstandardized Coefficients		Standardized Coefficients	t	Sig.	Collinearity Statistics	
		B	Std. Error	Beta			Tolerance	VIF
1	(Constant)	8.219	.696		11.810	.000		
	BOPO	-.096	.009	-.852	-11.280	.000	.786	1.272
	NIM	.270	.023	.877	11.788	.000	.809	1.236
	Inflasi	.010	.010	.073	1.062	.296	.949	1.054


a. Dependent Variable: ROA

Charts

Normal P-P Plot of Regression Standardized Residual



Scatterplot



```
NPART TESTS  
/K-S (NORMAL)=RES_1  
/MISSING ANALYSIS.
```

NPar Tests

[DataSet0]

One-Sample Kolmogorov-Smirnov Test

		Unstandardized Residual
N		38
Normal Parameters ^a	Mean	.0000000
	Std. Deviation	.19886209
Most Extreme Differences	Absolute	.139
	Positive	.105
	Negative	-.139
Kolmogorov-Smirnov Z		.854
Asymp. Sig. (2-tailed)		.459
a. Test distribution is Normal.		

Descriptives

Descriptive Statistics

	N	Minimum	Maximum	Mean	Std. Deviation
ROA	38	.16	2.02	.9368	.50959
BOPO	38	78.07	98.60	86.3177	4.49966
NIM	38	1.30	6.74	3.6062	1.65524
Inflasi	38	.17	15.74	6.6542	3.60843
Valid N (listwise)	38				