

DAFTAR PUSTAKA

- ACOG (2014) ‘Obstetric Data Definitions version 1.0’, *Revitalize*, 1(1), pp. 1–5.
- ACOG (2016) ‘Weight Gain During Pregnancy’, *The American College of Obstetricians and Gynecologists*, 548(548), pp. 1–3. doi: 10.1002/14651858.CD001877.
- Adhi, Y. B. (2010) *Ukuran lingkar lengan atas (Lila) ibu hamil dan kejadian Bayi Berat Badan Lahir Rendah (BBLR)*. UNS.
- Agrawal, S. and Singh, A. (2016) ‘Obesity or Underweight — What is Worse in Pregnancy ?’, *The Journal of Obstetrics and Gynecology of India*. Springer India, 66(6), pp. 448–452. doi: 10.1007/s13224-015-0735-4.
- Akter, S. M. et al. (2012) ‘Effects of third trimester counseling on pregnancy weight gain , birthweight , and breastfeeding among urban poor women in Bangladesh’, 33(3), pp. 194–201.
- Al-Shaikh, G. K. et al. (2017) ‘Grand multiparity and the possible risk of adverse maternal and neonatal outcomes: A dilemma to be deciphered’, *BMC Pregnancy and Childbirth*. BMC Pregnancy and Childbirth, 17(1), pp. 1–7. doi: 10.1186/s12884-017-1508-0.
- Aliyu, M. H. et al. (2005) ‘High Parity and Fetal Morbidity Outcomes’, 105(5), pp. 1045–1051. doi: 10.1097/01.AOG.0000157444.74674.75.
- Aminian, O. et al. (2014) ‘gestational age Association between maternal work activity on birth weight and’, 3(3), pp. 200–203. doi: 10.1016/S2305-0500(14)60026-9.
- Arshad, R., Karim, N. and Hasan, J. A. (2014) ‘Effects of insulin on placental, fetal and maternal outcomes in gestational diabetes mellitus’, *Pakistan Journal of Medical Sciences*, 30(2), pp. 240–244. doi: 10.12669/pjms.302.4396.
- Assefa, N., Berhane, Y. and Worku, A. (2012) ‘Wealth status, mid upper arm circumference (MUAC) and Ante Natal Care (ANC) are determinants for low birth weight in Kersa, Ethiopia’, *PLoS ONE*, 7(6). doi: 10.1371/journal.pone.0039957.
- Bener, A. et al. (2012) ‘Pattern of Maternal Complications and Low Birth Weight : Associated Risk Factors among Highly Endogamous Women’, 2012. doi: 10.5402/2012/540495.
- Borodulin, K., Evenson, K. R. and Herring, A. H. (2009) ‘Physical activity patterns during pregnancy through postpartum’, *BMC Women’s Health*, 9, pp. 1–7. doi: 10.1186/1472-6874-9-32.
- BPS Kabupaten Pacitan (2018) *Kabupaten Pacitan Dalam Angka*. Pacitan: BPS Kabupaten Pacitan.
- BPS Kota Tangerang Selatan (2015) *Kota Tangerang Selatan dalam Angka*.
- BPS Provinsi Jawa Timur (2018) *Provinsi Jawa Timur Dalam Angka*. BPS Provinsi Jawa Timur. Available at: <https://jatim.bps.go.id/>.

- Brito, N. B. *et al.* (2016) 'Relationship between Mid-Upper Arm Circumference and Body Mass Index in Inpatients', 78(cm), pp. 1–10. doi: 10.1371/journal.pone.0160480.
- Budiman, A. R. (2013) *Pengetahuan dan Sikap Penelitian Kesehatan*. Jakarta: Salemba Medika.
- Cahyani, W. (2015) *Hubungan antara Tingkat Pendidikan, Umur, Paritas, dan Kadar Hemoglobin pada Maternal dengan Kejadian BBLR di RS. dr.Oen Surakarta Tahun 2014*. UMS.
- Candrasari, A. *et al.* (2012) 'HUBUNGAN ANTARA PERTAMBAHAN BERAT BADAN IBU HAMIL DENGAN BERAT BADAN LAHIR BAYI DI KABUPATEN SEMARANG', *Jurnal Penelitian Sains & Teknologi*, 13(1), pp. 56–62.
- Cavazos-rehg, P. A. *et al.* (2016) 'Maternal age and risk of labor and delivery complications', *Maternal and Child Health Journal*, 19(6), pp. 1202–1211. doi: 10.1007/s10995-014-1624-7.Maternal.
- Cheung, T.-F. (2000) 'Pregnancy Weight Management. Adam-Media: Canada', *Adam-media : Canada*.
- Coo, H. *et al.* (2017) 'Interpregnancy Interval and Adverse Perinatal Outcomes : A Record-Linkage Study Using the Manitoba Population Research Data Repository', *JOGC*. Elsevier Inc, 39(6), pp. 420–433. doi: 10.1016/j.jogc.2017.01.010.
- Cunningham, F. G. *et al.* (2014) *William Obstetric*. 24th edn, *Mc Graw Hill Education*. 24th edn. Mc Graw Hill. doi: 10.13140/2.1.1814.6566.
- Darmayanti, Wilopo, S. A. and Nurdianti, D. S. (2010) 'Pengaruh Kenaikan Berat Badan Rata-Rata Per Minggu Pada Kehamilan Trimester II dan Trimester III Terhadap Risiko Berat Bayi Lahir Rendah', *Jurnal Berita Kedokteran Masyarakat*, 26(1), pp. 40–46.
- Depkes RI (2006) *Ibu Sehat Bayi Sehat*. Jakarta.
- Dewey, K. G. and Cohen, R. J. (2007) 'Does birth spacing affect maternal or child nutritional status ? A systematic literature review', pp. 151–173.
- Dewi, astrid purna (2009) 'hubungan antara kenaikan berat badan ibu hamil dengan kejadian bayi berat lahir rendah (BBLR) Di RSUD Dr. MOEWARDI SURAKARTA', 39, p. 12.
- Drehmer, M. *et al.* (2013) 'Association of Second and Third Trimester Weight Gain in Pregnancy with Maternal and Fetal Outcomes', *Plos One*, 8(1). doi: 10.1371/journal.pone.0054704.
- Durie, D. E. (2011) 'Effect of Second-Trimester and Third-Trimester Rate of Gestational Weight Gain on Maternal and Neonatal Outcomes', 118(3), pp. 18–20. doi: 10.1097/AOG.0b013e3182289f42.
- Endalamaw, A. *et al.* (2018) 'Low birth weight and its associated factors in Ethiopia: a systematic review and meta-analysis', *Italian Journal of Pediatrics*.

Italian Journal of Pediatrics, 44(1), pp. 1–12. doi: 10.1186/s13052-018-0586-6.

Fajriana, A. and Buanasita, A. (2016) ‘Faktor risiko yang berhubungan dengan kejadian bayi berat lahir rendah di kecamatan semampir surabaya’, (2018), pp. 71–80. doi: 10.20473/mgi.v13i1.71.

Frederick, I. O. *et al.* (2008) ‘Pre-pregnancy Body Mass Index , Gestational Weight Gain , and Other Maternal Characteristics in Relation to Infant Birth Weight’, pp. 557–567. doi: 10.1007/s10995-007-0276-2.

Hack, M. *et al.* (2011) ‘CHANGE IN PREVALENCE OF CHRONIC CONDITIONS BETWEEN CHILDHOOD AND ADOLESCENCE AMONG EXTREMELY LOW BIRTH WEIGHT CHILDREN’, 306(4), pp. 394–401. doi: 10.1001/jama.2011.1025.CHANGE.

Hanretty, K. P. (2010) *Obstetrics Illustrated*. 7th edn. Glasgow: Elsevier.

Hasan, T., Khan, A. and Ahmed, T. (2019) ‘Inadequate maternal weight gain in the third trimester increases the risk of intrauterine growth restriction in rural Bangladesh’, *PLoS ONE*, 8, pp. 1–14.

IOM and National Research Council (NRC) (2010) ‘Implementing Guidelines on Weight Gain Pregnancy’.

Jayanti, F. A., Dharmawan, Y. and Aruben, R. (2017) ‘Faktor-Faktor yang Berhubungan dengan Kejadian Berat Badan Lahir Rendah di Wilayah Kerja Puskesmas Bangetayu Kota Semarang Tahun 2016’, *Jurnal Kesehatan Masyarakat*, 5(4), pp. 812–822.

Kasim, F., Surachman, T. and Ruswandini (2011) ‘Hubungan antara Karakteristik Ibu Hamil dengan Kejadian Bayi Berat Badan Lahir Rendah di Rumah Sakit Immanuel Bandung Tahun 2008’, *JKM*, 10(2).

Kemenkes (2014) ‘Rencana Strategis Kementerian Kesehatan Republik Indonesia Tahun 2015-2019’, *Pusat Komunikasi Publik*. doi: 351.077 Ind r.

Kemenkes (2015) ‘InfoDATIN’, pp. 1–8.

Kemenkes RI (2010a) ‘Pedoman Pelayanan Antenatal’, *Pedoman Pelayanan Antenatal Terpadu*, p. 1 of 98. doi: 10.1016/j.bbr.2012.09.037.

Kemenkes RI (2010b) ‘Pedoman Pemantauan Wilayah Setempat Kesehatan Ibu dan Anak’, *Kementrian Kesehatan*, p. 1 of 76.

Kemenkes RI (2011) *Buku Acuan Manajemen Bayi Berat Lahir Rendah (BBLR) Untuk Bidan di Desa*. Jakarta.

Kemenkes RI (2015) *Infodatin: Situasi Kesehatan Reproduksi Remaja*. Jakarta: Kemenkes.

Khojasteh, F., Arbabisarjou, A. and Boryri, T. (2016) ‘The Relationship between Maternal Employment Status and Pregnancy Outcomes’, 8(9), pp. 37–43. doi: 10.5539/gjhs.v8n9p37.

Krishnan, M. (2014) ‘Profil Kelahiran Bayi Dengan Berat Badan Lahir Rendah (BBLR) di RSUP Haji Adam Malik Medan Tahun 2011-2012’, *USU*.

- Krukowski, R. A. *et al.* (2016) ‘Are early first trimester weights valid proxies for preconception weight?’, *BMC Pregnancy and Childbirth*. BMC Pregnancy and Childbirth, pp. 10–15. doi: 10.1186/s12884-016-1159-6.
- Kusumaningrum, A. I. (2012) ‘Hubungan Faktor Ibu dengan Kejadian Bayi Berat Lahir Rendah (BBLR) di Wilayah Kerja Puskesmas Gemawang Kecamatan Gemawang Kabupaten Temanggung Provinsi Temanggung’, *FKM UI*.
- Low, P. *et al.* (2005) ‘Factors affecting antenatal care attendance by mothers of Pacific infants living in New Zealand.’, *The New Zealand medical journal*, 118(1216).
- Macdonald-Wallis, C. *et al.* (2014) ‘Associations of blood pressure change in pregnancy with fetal growth and gestational age at delivery: Findings from a prospective cohort’, *Hypertension*, 64(1), pp. 36–44. doi: 10.1161/HYPERTENSIONAHA.113.02766.
- Maghfiroh, L. (2015) *Pertambahan Berat Badan Ibu Hamil dan Kejadian Berat Bayi Lahir Rendah (BBLR) di Wilayah Kerja Puskesmas Pamulang Kota Tangerang Selatan Tahun 2013-2015*, UIN Syarif Hidayatullah. doi: 10.1017/CBO9781107415324.004.
- Mahmoodi, Z. *et al.* (2016) ‘Association of maternal working condition with low birth weight: The social determinants of health approach’, *Annals of Medical and Health Sciences Research*, 5(6), p. 385. doi: 10.4103/2141-9248.177982.
- Maindarker, G. *et al.* (2012) ‘Maternal risk factors associated with term low birth weight neonates: A matched-pair case control study’, *Indian Pediatrics*, 49(1), pp. 25–28.
- Mala, T. M. (1988) ‘Interpregnancy interval and low birth weight: findings from a case-control study’, 128(5), pp. 1111–1116.
- Manisrilyati, R. (2012) ‘Hubungan Paritas Ibu Bersalin dengan Kejadian Bayi Berat Lahir Rendah di Puskesmas Mergangsan Kota Yogyakarta Tahun 2010’.
- Manuaba, I. B. (2000) *Kapita Selekta Penatalaksanaan Rutin Obstetri Ginekologi dan KB*. Jakarta: EGC.
- Merklinger-Gruchala, A., Jasienska, G. and Kapiszewska, M. (2015) ‘Short Interpregnancy Interval and Low Birth Weight: A Role of Parity’, *American Journal of Biology*, 00(January), pp. 1–7. doi: 10.1002/ajhb.22708.
- Mubarak, W. I. *et al.* (2007) *Promosi Kesehatan : Sebuah Pengantar Proses Belajar Mengajar dalam Pendidikan*. Yogyakarta: Graha Ilmu.
- Nasreen, H. E. *et al.* (2010) ‘Low birth weight in offspring of women with depressive and anxiety symptoms during pregnancy: Results from a population based study in Bangladesh’, *BMC Public Health*, 10. doi: 10.1186/1471-2458-10-515.
- Niedhammer, I. *et al.* (2009) ‘Occupational predictors of pregnancy outcomes in Irish working women in the Lifeways cohort’, *BJOG: An International Journal of Obstetrics and Gynaecology*, 116(7), pp. 943–952. doi: 10.1111/j.1471-0528.2009.02160.x.

- Nyaruhucha, C. N. M. *et al.* (2006) ‘Maternal weight gain in second and third trimesters and their relationship with birth weights in Morogoro Municipality , Tanzania’, 8(1), pp. 41–44.
- O’Leary, M. *et al.* (2017) ‘A cohort study of low birth weight and health outcomes in the first year of life, Ghana’, *Bulletin of the World Health Organization*, 95(8), pp. 574–583. doi: 10.2471/BLT.16.180273.
- Pathirathna, M. L. *et al.* (2019) ‘Effects of Physical Activity During Pregnancy on Neonatal Birth’, pp. 1–8. doi: 10.1038/s41598-019-42473-7.
- Petry, C. J. *et al.* (2018) ‘Vomiting in pregnancy is associated with a higher risk of low birth weight : a cohort study’. *BMC Pregnancy and Childbirth*, pp. 1–8.
- Pinontoan, V. M. and Tombokan, S. G. . (2015) ‘Hubungan Umur dan Paritas dengan Kejadian Bayi Berat Lahir Rendah’, *Jurnal Ilmiah Bidan*, 3(1), pp. 20–25.
- Pramono, M. S. and Putro, G. (2009) ‘Risiko Terjadinya Berat Bayi Lahir Rendah Menurut Determinan Sosial, Ekonomi, dan Demografi di Indonesia’, *Buletin Penelitian Sistem Kesehatan*, 12(2).
- Promkes Kemenkes (2018) *Pentingnya Pemeriksaan Kehamilan (ANC) di Fasilitas Kesehatan, Direktorat Promosi Kesejahteraan dan Pemberdayaan Masyarakat Kementerian Kesehatan*. Available at: <http://promkes.kemkes.go.id/pentingnya-pemeriksaan-kehamilan-anc-di-fasilitas-kesehatan> (Accessed: 8 April 2019).
- Rahman, L. A., Hairi, N. N. and Salleh, N. (2008) ‘Association Between Pregnancy Induced Hypertension and Low Birth Weight; A Population Based Case-Control Study’, *Asia Pacific Journal of Public Health*, 20(2), pp. 152–158. doi: 10.1177/1010539507311553.
- Rasmussen, K. M., Catalano, P. M. and Yaktine, A. L. (2009) ‘New guidelines for weight gain during pregnancy: what obstetrician/gynecologists should know’, *Current Opinion in Obstetrics and Gynecology*, 21(6), pp. 521–526. doi: 10.1097/gco.0b013e328332d24e.
- Ravenalla (2015) *Hubungan Antenatal Care (ANC) dengan Kejadian Bayi Berat Lahir Rendah di Kabupaten Purbalingga*. UGM. Available at: http://etd.repository.ugm.ac.id/index.php?mod=penelitian_detail&sub=Penelitian Detail&act=view&typ=html&buku_id=84948&obyek_id=4.
- Restrepo-méndez, M. C. *et al.* (2015) ‘The Association of Maternal Age with Birthweight and Gestational Age : A Cross-Cohort Comparison’, pp. 31–40. doi: 10.1111/ppe.12162.
- Reyes, L. and Manalich, R. (2015) *Long-term consequences of low birth weight*. Cuba: Instituto de Nefrología. doi: 10.1111/j.1523-1755.2005.09718.x.
- Reyes, L. and Mañalich, R. (2005) ‘Long-term consequences of low birth weight’, *Kidney International, Supplement*, 68(97), pp. 107–111. doi: 10.1111/j.1523-1755.2005.09718.x.
- Roth, J. *et al.* (1998) ‘The Risk of Teen Mothers having Low Birth Weight Babies:Implication of Recent Medical Research for School Health Personnel’,

Journal of School Health, 68(7), pp. 271–275.

Seely, E. W. and Ecker, J. (2014) ‘Chronic hypertension in pregnancy’, *Circulation*, 129(11), pp. 1254–1261. doi: 10.1161/CIRCULATIONAHA.113.003904.

Septiani, R. (2015) ‘Faktor Maternal Pada Kejadian Berat Badan Lahir Rendah (BBLR) Di Indonesia (Analisis Data Riskesdas 2013)’, *UIN Syarif Hidayatullah*. Jakarta. Available at: <http://repository.uinjkt.ac.id/dspace/bitstream/123456789/28928/1/Rini Septiani-fkik.pdf>.

Shabila, N. P., Ahmed, H. M. and Yasin, M. Y. (2015) ‘Women ’ s views and experiences of antenatal care in Iraq : a Q methodology study’.

Shah, P. S. (2010) ‘Parity and low birth weight and preterm birth: a systematic review and meta-analyses.’, *Acta Obstetricia et Gynecologica Scandinavica*. doi: 10.3109/00016349.2010.486827.

Shah, P. S. *et al.* (2011) ‘Intention to Become Pregnant and Low Birth Weight and Preterm Birth: A Systematic Review’, *Maternal and Child Health Journal*, 15(2), pp. 205–216.

Soetjiningsih (1995) *Tumbuh Kembang Anak*. Edited by G. Ranuh. Jakarta: EGC.

Solang, S., Purwandari, A. and Lohoraung, A. (2012) ‘Hubungan Kepuasan Pelayanan Antenatal Care (ANC) dengan Frekuensi Kunjungan Ibu Hamil di Puskesmas Kota Kombos Kecamat Singkil Kota Manado’, *GIZIDO*, 4(1).

Soma-pillay, P. *et al.* (2016) ‘Physiological changes in pregnancy’, 27(2), pp. 89–94. doi: 10.5830/CVJA-2016-021.

Tazkiah, M. (2013) *Determinan Epidemiologi Kejadian BBLR pada Daerah Endemis Malaria di Kabupaten Banjar Provinsi Kalimantan Selatan*. UNAIR.

Ullah, M. A. *et al.* (2003) ‘Biological Risk Factors of Low Birth Weight in Rural Rajshahi’, 16(2), pp. 2–5.

Vaswani, P. R. and Sabharwal, S. (2013) ‘Trends in the Occurrence of Antenatal and Perinatal Complications with Increasing Parity’, *Journal of Obstetrics and Gynecology of India*, 63(4), pp. 260–267. doi: 10.1007/s13224-012-0344-4.

Virdaus, S., Hakimi, M. and Berty Murtiningsih (2011) ‘Kurang Energi Kronis Ibu Hamil sebagai Faktor Risiko Bayi Berat Lahir Rendah’, *Berita Kedokteran Masyarakat*, 27(4), pp. 187–196. Available at: <http://berita-kedokteran-masyarakat.org/index.php/BKM/article/view/291>.

Watson, E. D. *et al.* (2018) ‘The Influence of Objectively Measured Physical Activity During Pregnancy on Maternal and Birth Outcomes in Urban Black South African Women’, *Maternal and Child Health Journal*. Springer US, 0(0), p. 0. doi: 10.1007/s10995-018-2504-3.

WHO (2014) ‘Global Nutrition Targets 2025 Low Birth Weight Policy Brief’. Switzerland: WHO, p. 8.

Yang, W. *et al.* (2017) ‘Relationship Between Gestational Weight Gain and

Pregnancy Complications or Delivery Outcome', *Scientific Reports*. Springer US, (11), pp. 1–9. doi: 10.1038/s41598-017-12921-3.

Yongky *et al.* (2009) 'Status Gizi Awal Kehamilan dan Pertambahan Berat Badan Ibu Hamil Kaitannya dengan BBLR', *Jurnal Gizi dan Pangan*, 1(41), pp. 8–12. doi: <http://dx.doi.org/10.25182/jgp.2009.4.1.8-12>.

Zanardo, V. *et al.* (2016) 'Gestational weight gain and fetal growth in underweight women', *Italian Journal of Pediatrics*. Italian Journal of Pediatrics, 42(1), pp. 1–6. doi: 10.1186/s13052-016-0284-1.

