

#### Lampiran 4. Hasil SPSS distribusi karakteristik demografi dan klinis responden

### Frequencies

#### Statistics

usia

N	Valid	100
	Missing	0

usia

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	kurang dari sama dengan 50 tahun	35	35.0	35.0	35.0
	lebih dari 50 tahun	65	65.0	65.0	100.0
	Total	100	100.0	100.0	

### Frequencies

#### Statistics

JenisKelamin

N	Valid	100
	Missing	0

JenisKelamin

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	pria	33	33.0	33.0	33.0
	wanita	67	67.0	67.0	100.0
	Total	100	100.0	100.0	

## Frequencies

### Statistics

JumlahAntihipertensi

N	Valid	100
	Missing	0

### JumlahAntihipertensi

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	tunggal	49	49.0	49.0	49.0
	kombinasi	51	51.0	51.0	100.0
	Total	100	100.0	100.0	

## Frequencies

### Statistics

JumlahPenyakitPenyerta

N	Valid	100
	Missing	0

### JumlahPenyakitPenyerta

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	0	71	71.0	71.0	71.0
	1+	29	29.0	29.0	100.0
	Total	100	100.0	100.0	

## Frequencies

### Statistics

pendidikan

N	Valid	100
	Missing	0

### pendidikan

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	kurang dari sama dengan SMP	62	62.0	62.0	62.0
	lebih dari SMP	38	38.0	38.0	100.0
Total		100	100.0	100.0	

## Frequencies

### Statistics

pekerjaan

N	Valid	100
	Missing	0

### pekerjaan

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	formal	27	27.0	27.0	27.0
	non formal	73	73.0	73.0	100.0
Total		100	100.0	100.0	

## Frequencies

### Statistics

kepatuhan

N	Valid	100
	Missing	0

### kepatuhan

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	patuh	43	43.0	43.0	43.0
	tidak patuh	57	57.0	57.0	100.0
Total		100	100.0	100.0	

## Frequencies

### Statistics

pengetahuan

N	Valid	100
	Missing	0

### pengetahuan

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	tinggi	49	49.0	49.0	49.0
	rata-rata	31	31.0	31.0	80.0
	rendah	20	20.0	20.0	100.0
Total		100	100.0	100.0	

## Frequencies

### Statistics

		SpecificNecessity	SpesificConcerns	GeneralHarmful	GeneralOveruse
N	Valid	100	100	100	100
	Missing	0	0	0	0

## Frequency Table

**SpecificNecessity**

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	tinggi	76	76.0	76.0	76.0
	rendah	24	24.0	24.0	100.0
	Total	100	100.0	100.0	

**SpecificConcerns**

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	tinggi	60	60.0	60.0	60.0
	rendah	40	40.0	40.0	100.0
	Total	100	100.0	100.0	

**GeneralHarmful**

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	tinggi	87	87.0	87.0	87.0
	rendah	13	13.0	13.0	100.0
	Total	100	100.0	100.0	

**GeneralOveruse**

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	tinggi	55	55.0	55.0	55.0
	rendah	45	45.0	45.0	100.0
	Total	100	100.0	100.0	

### Lampiran 5. Hasil SPSS Tingkat Pengetahuan Pasien Hipertensi

#### 1. Tingkat pengetahuan berdasarkan usia

##### Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
usia * pengetahuan	100	100.0%	0	.0%	100	100.0%

##### usia \* pengetahuan Crosstabulation

Count		pengetahuan			Total
		tinggi	rata-rata	rendah	
usia	kurang dari sama dengan 50 tahun	22	11	2	35
	lebih dari 50 tahun	27	20	18	65
Total		49	31	20	100

##### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	7.608 <sup>a</sup>	2	.022
Likelihood Ratio	8.744	2	.013
Linear-by-Linear Association	6.968	1	.008
N of Valid Cases	100		

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 7.00.

## 2. Tingkat pengetahuan berdasarkan jenis kelamin

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
JenisKelamin * pengetahuan	100	100.0%	0	.0%	100	100.0%

JenisKelamin \* pengetahuan Crosstabulation

Count		pengetahuan			
		tinggi	rata-rata	rendah	Total
		JenisKelamin	pria	18	9
	wanita	31	22	14	67
Total		49	31	20	100

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.611 <sup>a</sup>	2	.737
Likelihood Ratio	.612	2	.736
Linear-by-Linear Association	.436	1	.509
N of Valid Cases	100		

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 6.60.

## 3. Tingkat Pengetahuan berdasarkan jumlah antihipertensi

**Case Processing Summary**

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
JumlahAntihipertensi * pengetahuan	100	100.0%	0	.0%	100	100.0%

**JumlahAntihipertensi \* pengetahuan Crosstabulation**

Count		pengetahuan			Total
		tinggi	rata-rata	rendah	
		JumlahAntihipertensi tunggal	30	10	
	JumlahAntihipertensi kombinasi	19	21	11	51
	Total	49	31	20	100

**Chi-Square Tests**

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	6.535 <sup>a</sup>	2	.038
Likelihood Ratio	6.640	2	.036
Linear-by-Linear Association	3.014	1	.083
N of Valid Cases	100		

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 9.80.



## 4. Tingkat pengetahuan berdasarkan jumlah penyakit penyerta

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
JumlahPenyakitPenyerta * pengetahuan	100	100.0%	0	.0%	100	100.0%

JumlahPenyakitPenyerta \* pengetahuan Crosstabulation

Count		pengetahuan			Total
		tinggi	rata-rata	rendah	
		JumlahPenyakitPenyerta	0	31	
	1+	18	6	5	29
Total		49	31	20	100

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	2.980 <sup>a</sup>	2	.225
Likelihood Ratio	3.037	2	.219
Linear-by-Linear Association	1.672	1	.196
N of Valid Cases	100		

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 5.80.

## 5. Tingkat pengetahuan berdasarkan pendidikan

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
pendidikan * pengetahuan	100	100.0%	0	.0%	100	100.0%

pendidikan \* pengetahuan Crosstabulation

Count		pengetahuan			Total
		tinggi	rata-rata	rendah	
		pendidikan kurang dari sama dengan SMP	20	23	
lebih dari SMP	29	8	1	38	
Total		49	31	20	100

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	20.534 <sup>a</sup>	2	.000
Likelihood Ratio	23.203	2	.000
Linear-by-Linear Association	19.996	1	.000
N of Valid Cases	100		

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 7.60.

## 6. Tingkat pengetahuan berdasarkan pekerjaan

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
pekerjaan * pengetahuan	100	100.0%	0	.0%	100	100.0%

pekerjaan \* pengetahuan Crosstabulation

Count		pengetahuan			Total
		tinggi	rata-rata	rendah	
pekerjaan	formal	19	6	2	27
	non formal	30	25	18	73
Total		49	31	20	100

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	7.299 <sup>a</sup>	2	.026
Likelihood Ratio	7.748	2	.021
Linear-by-Linear Association	6.971	1	.008
N of Valid Cases	100		

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 5.40.

### Lampiran 6. Hasil SPSS Tingkat Kepercayaan Pasien Hipertensi

#### 1. Tingkat kepercayaan berdasarkan usia

##### a. Tingkat kepercayaan tentang kebutuhan obat

#### Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
usia * SpecificNecessity	100	100.0%	0	.0%	100	100.0%

#### usia \* SpecificNecessity Crosstabulation

Count		SpecificNecessity		Total
		tinggi	rendah	
usia	kurang dari sama dengan 50 tahun	28	7	35
	lebih dari 50 tahun	48	17	65
Total		76	24	100

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.472 <sup>a</sup>	1	.492		
Continuity Correction <sup>b</sup>	.195	1	.659		
Likelihood Ratio	.482	1	.488		
Fisher's Exact Test				.625	.334
Linear-by-Linear Association	.468	1	.494		
N of Valid Cases <sup>b</sup>	100				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 8.40.

b. Computed only for a 2x2 table

## b. Tingkat kepercayaan tentang kekhawatiran pada obat

**Case Processing Summary**

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
usia * SpesificConcerns	100	100.0%	0	.0%	100	100.0%

**usia \* SpesificConcerns Crosstabulation**

Count		SpesificConcerns		Total
		tinggi	rendah	
usia	kurang dari sama dengan 50 tahun	14	21	35
	lebih dari 50 tahun	46	19	65
Total		60	40	100

**Chi-Square Tests**

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	8.974 <sup>a</sup>	1	.003		
Continuity Correction <sup>b</sup>	7.738	1	.005		
Likelihood Ratio	8.945	1	.003		
Fisher's Exact Test				.005	.003
Linear-by-Linear Association	8.885	1	.003		
N of Valid Cases <sup>b</sup>	100				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 14.00.

b. Computed only for a 2x2 table

## c. Tingkat kepercayaan tentang penggunaan obat berlebih

**Case Processing Summary**

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
usia * GeneralOveruse	100	100.0%	0	.0%	100	100.0%

**usia \* GeneralOveruse Crosstabulation**

Count		GeneralOveruse		Total
		tinggi	rendah	
usia	kurang dari sama dengan 50 tahun	15	20	35
	lebih dari 50 tahun	40	25	65
Total		55	45	100

**Chi-Square Tests**

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	3.208 <sup>a</sup>	1	.073		
Continuity Correction <sup>b</sup>	2.498	1	.114		
Likelihood Ratio	3.208	1	.073		
Fisher's Exact Test				.093	.057
Linear-by-Linear Association	3.176	1	.075		
N of Valid Cases <sup>b</sup>	100				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 15.75.

b. Computed only for a 2x2 table

## d. Tingkat kepercayaan tentang bahaya obat

**Case Processing Summary**

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
usia * GeneralHarmful	100	100.0%	0	.0%	100	100.0%

**usia \* GeneralHarmful Crosstabulation**

Count		GeneralHarmful		Total
		tinggi	rendah	
usia	kurang dari sama dengan 50 tahun	29	6	35
	lebih dari 50 tahun	58	7	65
Total		87	13	100

**Chi-Square Tests**

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.817 <sup>a</sup>	1	.366		
Continuity Correction <sup>b</sup>	.351	1	.554		
Likelihood Ratio	.791	1	.374		
Fisher's Exact Test				.369	.272
Linear-by-Linear Association	.809	1	.368		
N of Valid Cases <sup>b</sup>	100				

a. 1 cells (25.0%) have expected count less than 5. The minimum expected count is 4.55.

b. Computed only for a 2x2 table

2. Tingkat kepercayaan berdasarkan jenis kelamin  
 a. Tingkat kepercayaan tentang kebutuhan obat

#### Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
JenisKelamin * SpecificNecessity	100	100.0%	0	.0%	100	100.0%

#### JenisKelamin \* SpecificNecessity Crosstabulation

Count		SpecificNecessity		Total
		tinggi	rendah	
JenisKelamin	pria	24	9	33
	wanita	52	15	67
Total		76	24	100

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.289 <sup>a</sup>	1	.591		
Continuity Correction <sup>b</sup>	.083	1	.773		
Likelihood Ratio	.285	1	.593		
Fisher's Exact Test				.624	.381
Linear-by-Linear Association	.286	1	.593		
N of Valid Cases <sup>b</sup>	100				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 7.92.

b. Computed only for a 2x2 table



## b. Tingkat kepercayaan tentang kekhawatiran pada obat

**Case Processing Summary**

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
JenisKelamin * SpesificConcerns	100	100.0%	0	.0%	100	100.0%

**JenisKelamin \* SpesificConcerns Crosstabulation**

Count		SpesificConcerns		
		tinggi	rendah	Total
JenisKelamin	pria	21	12	33
	wanita	39	28	67
Total		60	40	100

**Chi-Square Tests**

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.271 <sup>a</sup>	1	.602		
Continuity Correction <sup>b</sup>	.092	1	.761		
Likelihood Ratio	.273	1	.601		
Fisher's Exact Test				.668	.383
Linear-by-Linear Association	.269	1	.604		
N of Valid Cases <sup>b</sup>	100				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 13.20.

b. Computed only for a 2x2 table

## c. Tingkat kepercayaan tentang penggunaan obat berlebih

**Case Processing Summary**

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
JenisKelamin * GeneralOveruse	100	100.0%	0	.0%	100	100.0%

**JenisKelamin \* GeneralOveruse Crosstabulation**

Count		GeneralOveruse		
		tinggi	rendah	Total
JenisKelamin	pria	15	18	33
	wanita	40	27	67
Total		55	45	100

**Chi-Square Tests**

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	1.813 <sup>a</sup>	1	.178		
Continuity Correction <sup>b</sup>	1.283	1	.257		
Likelihood Ratio	1.810	1	.179		
Fisher's Exact Test				.204	.129
Linear-by-Linear Association	1.795	1	.180		
N of Valid Cases <sup>b</sup>	100				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 14.85.

b. Computed only for a 2x2 table

## d. Tingkat kepercayaan tentang bahaya obat

**Case Processing Summary**

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
JenisKelamin * GeneralHarmful	100	100.0%	0	.0%	100	100.0%

**JenisKelamin \* GeneralHarmful Crosstabulation**

Count		GeneralHarmful		Total
		tinggi	rendah	
JenisKelamin	pria	30	3	33
	wanita	57	10	67
Total		87	13	100

**Chi-Square Tests**

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.665 <sup>a</sup>	1	.415		
Continuity Correction <sup>b</sup>	.250	1	.617		
Likelihood Ratio	.702	1	.402		
Fisher's Exact Test				.536	.317
Linear-by-Linear Association	.659	1	.417		
N of Valid Cases <sup>b</sup>	100				

a. 1 cells (25.0%) have expected count less than 5. The minimum expected count is 4.29.

b. Computed only for a 2x2 table

## 3. Tingkat kepercayaan berdasarkan jumlah antihipertensi

## a. Tingkat kepercayaan tentang kebutuhan obat

**Case Processing Summary**

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
JumlahAntihipertensi * SpecificNecessity	100	100.0%	0	.0%	100	100.0%

**JumlahAntihipertensi \* SpecificNecessity Crosstabulation**

Count				
		SpecificNecessity		
		tinggi	rendah	Total
JumlahAntihipertensi	tunggal	34	15	49
	kombinasi	42	9	51
Total		76	24	100

**Chi-Square Tests**

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	2.303 <sup>a</sup>	1	.129		
Continuity Correction <sup>b</sup>	1.647	1	.199		
Likelihood Ratio	2.320	1	.128		
Fisher's Exact Test				.162	.100
Linear-by-Linear Association	2.280	1	.131		
N of Valid Cases <sup>b</sup>	100				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 11.76.

b. Computed only for a 2x2 table

## b. Tingkat kepercayaan tentang kekhawatiran pada obat

**Case Processing Summary**

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
JumlahAntihipertensi * SpesificConcerns	100	100.0%	0	.0%	100	100.0%

**JumlahAntihipertensi \* SpesificConcerns Crosstabulation**

Count				
		SpesificConcerns		Total
		tinggi	rendah	
JumlahAntihipertensi	tunggal	28	21	49
	kombinasi	32	19	51
Total		60	40	100

**Chi-Square Tests**

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.327 <sup>a</sup>	1	.568		
Continuity Correction <sup>b</sup>	.135	1	.713		
Likelihood Ratio	.327	1	.567		
Fisher's Exact Test				.684	.357
Linear-by-Linear Association	.324	1	.569		
N of Valid Cases <sup>b</sup>	100				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 19.60.

b. Computed only for a 2x2 table

## c. Tingkat kepercayaan tentang penggunaan obat berlebih

**Case Processing Summary**

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
JumlahAntihipertensi * GeneralOveruse	100	100.0%	0	.0%	100	100.0%

**JumlahAntihipertensi \* GeneralOveruse Crosstabulation**

Count		GeneralOveruse		
		tinggi	rendah	Total
JumlahAntihipertensi	tunggal	24	25	49
	kombinasi	31	20	51
Total		55	45	100

**Chi-Square Tests**

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	1.407 <sup>a</sup>	1	.236		
Continuity Correction <sup>b</sup>	.970	1	.325		
Likelihood Ratio	1.410	1	.235		
Fisher's Exact Test				.315	.162
Linear-by-Linear Association	1.393	1	.238		
N of Valid Cases <sup>b</sup>	100				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 22.05.

b. Computed only for a 2x2 table

## d. Tingkat kepercayaan tentang bahaya obat

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
JumlahAntihipertensi * GeneralHarmful	100	100.0%	0	.0%	100	100.0%

JumlahAntihipertensi \* GeneralHarmful Crosstabulation

Count				
		GeneralHarmful		Total
		tinggi	rendah	
JumlahAntihipertensi	tunggal	42	7	49
	kombinasi	45	6	51
Total		87	13	100

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.140 <sup>a</sup>	1	.708		
Continuity Correction <sup>b</sup>	.006	1	.938		
Likelihood Ratio	.140	1	.708		
Fisher's Exact Test				.772	.469
Linear-by-Linear Association	.139	1	.709		
N of Valid Cases <sup>b</sup>	100				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 6.37.

b. Computed only for a 2x2 table

## 4. Tingkat kepercayaan berdasarkan jumlah penyakit penyerta

## a. Tingkat kepercayaan tentang kebutuhan obat

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
JumlahPenyakitPenyerta * SpecificNecessity	100	100.0%	0	.0%	100	100.0%

JumlahPenyakitPenyerta \* SpecificNecessity Crosstabulation

Count		SpecificNecessity		Total
		tinggi	rendah	
JumlahPenyakitPenyerta	0	54	17	71
	1+	22	7	29
Total		76	24	100

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.000 <sup>a</sup>	1	.984		
Continuity Correction <sup>b</sup>	.000	1	1.000		
Likelihood Ratio	.000	1	.984		
Fisher's Exact Test				1.000	.586
Linear-by-Linear Association	.000	1	.984		
N of Valid Cases <sup>b</sup>	100				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 6.96.

b. Computed only for a 2x2 table



## b. Tingkat kepercayaan tentang kekhawatiran pada obat

**Case Processing Summary**

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
JumlahPenyakitPenyerta * SpescificConcerns	100	100.0%	0	.0%	100	100.0%

**JumlahPenyakitPenyerta \* SpescificConcerns Crosstabulation**

Count		SpescificConcerns		
		tinggi	rendah	Total
		JumlahPenyakitPenyerta	0	41
	1+	19	10	29
Total		60	40	100

**Chi-Square Tests**

	Value	df	Asymp. Sig. (2- sided)	Exact Sig. (2- sided)	Exact Sig. (1- sided)
Pearson Chi-Square	.518 <sup>a</sup>	1	.472		
Continuity Correction <sup>b</sup>	.245	1	.621		
Likelihood Ratio	.524	1	.469		
Fisher's Exact Test				.508	.312
Linear-by-Linear Association	.513	1	.474		
N of Valid Cases <sup>b</sup>	100				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 11.60.

b. Computed only for a 2x2 table

## c. Tingkat kepercayaan tentang penggunaan obat berlebih

**Case Processing Summary**

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
JumlahPenyakitPenyerta * GeneralOveruse	100	100.0%	0	.0%	100	100.0%

**JumlahPenyakitPenyerta \* GeneralOveruse Crosstabulation**

Count		GeneralOveruse		
		tinggi	rendah	Total
		JumlahPenyakitPenyerta	0	38
	1+	17	12	29
Total		55	45	100

**Chi-Square Tests**

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.216 <sup>a</sup>	1	.642		
Continuity Correction <sup>b</sup>	.059	1	.808		
Likelihood Ratio	.217	1	.641		
Fisher's Exact Test				.665	.405
Linear-by-Linear Association	.214	1	.644		
N of Valid Cases <sup>b</sup>	100				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 13.05.

b. Computed only for a 2x2 table

## d. Tingkat kepercayaan tentang bahaya obat

**Case Processing Summary**

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
JumlahPenyakitPenyerta * GeneralHarmful	100	100.0%	0	.0%	100	100.0%

**JumlahPenyakitPenyerta \* GeneralHarmful Crosstabulation**

Count		GeneralHarmful		
		tinggi	rendah	Total
		JumlahPenyakitPenyerta	0	61
	1+	26	3	29
Total		87	13	100

**Chi-Square Tests**

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.255 <sup>a</sup>	1	.614		
Continuity Correction <sup>b</sup>	.031	1	.860		
Likelihood Ratio	.265	1	.607		
Fisher's Exact Test				.751	.444
Linear-by-Linear Association	.252	1	.616		
N of Valid Cases <sup>b</sup>	100				

a. 1 cells (25.0%) have expected count less than 5. The minimum expected count is 3.77.

b. Computed only for a 2x2 table

## 5. Tingkat kepercayaan berdasarkan pendidikan

## a. tingkat kepercayaan tentang kebutuhan obat

**Case Processing Summary**

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
pendidikan * SpecificNecessity	100	100.0%	0	.0%	100	100.0%

**pendidikan \* SpecificNecessity Crosstabulation**

Count		SpecificNecessity		
		tinggi	rendah	Total
		pendidikan kurang dari sama dengan SMP	47	15
lebih dari SMP	29	9	38	
Total		76	24	100

**Chi-Square Tests**

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.003 <sup>a</sup>	1	.954		
Continuity Correction <sup>b</sup>	.000	1	1.000		
Likelihood Ratio	.003	1	.954		
Fisher's Exact Test				1.000	.576
Linear-by-Linear Association	.003	1	.954		
N of Valid Cases <sup>b</sup>	100				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 9.12.

b. Computed only for a 2x2 table

## b. tingkat kepercayaan tentang kekhawatiran pada obat

**Case Processing Summary**

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
pendidikan * SpescificConcerns	100	100.0%	0	.0%	100	100.0%

**pendidikan \* SpescificConcerns Crosstabulation**

Count		SpescificConcerns		
		tinggi	rendah	Total
		pendidikan kurang dari sama dengan SMP	41	21
lebih dari SMP	19	19	38	
Total		60	40	100

**Chi-Square Tests**

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	2.554 <sup>a</sup>	1	.110	.142	.083
Continuity Correction <sup>b</sup>	1.926	1	.165		
Likelihood Ratio	2.541	1	.111		
Fisher's Exact Test					
Linear-by-Linear Association	2.528	1	.112		
N of Valid Cases <sup>b</sup>	100				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 15.20.

b. Computed only for a 2x2 table

## c. Tingkat kepercayaan tentang penggunaan obat berlebih

**Case Processing Summary**

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
pendidikan * GeneralOveruse	100	100.0%	0	.0%	100	100.0%

**pendidikan \* GeneralOveruse Crosstabulation**

Count		GeneralOveruse		
		tinggi	rendah	Total
		pendidikan kurang dari sama dengan SMP	40	22
lebih dari SMP	15	23	38	
Total		55	45	100

**Chi-Square Tests**

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	5.970 <sup>a</sup>	1	.015		
Continuity Correction <sup>b</sup>	5.001	1	.025		
Likelihood Ratio	5.997	1	.014		
Fisher's Exact Test				.022	.013
Linear-by-Linear Association	5.910	1	.015		
N of Valid Cases <sup>b</sup>	100				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 17.10.

b. Computed only for a 2x2 table

## d. Tingkat kepercayaan tentang bahaya obat

## Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
pendidikan * GeneralHarmful	100	100.0%	0	.0%	100	100.0%

## pendidikan \* GeneralHarmful Crosstabulation

Count		GeneralHarmful		Total
		tinggi	rendah	
pendidikan	kurang dari sama dengan SMP	54	8	62
	lebih dari SMP	33	5	38
Total		87	13	100

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.001 <sup>a</sup>	1	.971	1.000	.599
Continuity Correction <sup>b</sup>	.000	1	1.000		
Likelihood Ratio	.001	1	.971		
Fisher's Exact Test					
Linear-by-Linear Association	.001	1	.971		
N of Valid Cases <sup>b</sup>	100				

a. 1 cells (25.0%) have expected count less than 5. The minimum expected count is 4.94.

b. Computed only for a 2x2 table

## 6. Tingkat kepercayaan berdasarkan pekerjaan

## a. Tingkat kepercayaan tentang kebutuhan

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
pekerjaan * SpecificNecessity	100	100.0%	0	.0%	100	100.0%

pekerjaan \* SpecificNecessity Crosstabulation

Count		SpecificNecessity		
		tinggi	rendah	Total
pekerjaan	formal	21	6	27
	non formal	55	18	73
Total		76	24	100

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.064 <sup>a</sup>	1	.800		
Continuity Correction <sup>b</sup>	.000	1	1.000		
Likelihood Ratio	.065	1	.799		
Fisher's Exact Test				1.000	.513
Linear-by-Linear Association	.063	1	.801		
N of Valid Cases <sup>b</sup>	100				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 6.48.

b. Computed only for a 2x2 table



## b. Tingkat kepercayaan tentang kekhawatiran pada obat

**Case Processing Summary**

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
pekerjaan * SpescificConcerns	100	100.0%	0	.0%	100	100.0%

**pekerjaan \* SpescificConcerns Crosstabulation**

Count		SpescificConcerns		
		tinggi	rendah	Total
pekerjaan	formal	13	14	27
	non formal	47	26	73
Total		60	40	100

**Chi-Square Tests**

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	2.165 <sup>a</sup>	1	.141		
Continuity Correction <sup>b</sup>	1.541	1	.214		
Likelihood Ratio	2.137	1	.144		
Fisher's Exact Test				.171	.108
Linear-by-Linear Association	2.143	1	.143		
N of Valid Cases <sup>b</sup>	100				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 10.80.

b. Computed only for a 2x2 table

c. tingkat kepercayaan tentang penggunaan obat berlebih

**Case Processing Summary**

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
pekerjaan * GeneralOveruse	100	100.0%	0	.0%	100	100.0%

**pekerjaan \* GeneralOveruse Crosstabulation**

Count		GeneralOveruse		
		tinggi	rendah	Total
pekerjaan	formal	10	17	27
	non formal	45	28	73
Total		55	45	100

**Chi-Square Tests**

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	4.822 <sup>a</sup>	1	.028		
Continuity Correction <sup>b</sup>	3.879	1	.049		
Likelihood Ratio	4.830	1	.028		
Fisher's Exact Test				.041	.024
Linear-by-Linear Association	4.774	1	.029		
N of Valid Cases <sup>b</sup>	100				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 12.15.

b. Computed only for a 2x2 table

## d. Tingkat kepercayaan tentang bahaya obat

## Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
pekerjaan * GeneralHarmful	100	100.0%	0	.0%	100	100.0%

## pekerjaan \* GeneralHarmful Crosstabulation

Count		GeneralHarmful		Total
		tinggi	rendah	
pekerjaan	formal	23	4	27
	non formal	64	9	73
Total		87	13	100

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.108 <sup>a</sup>	1	.743		
Continuity Correction <sup>b</sup>	.000	1	1.000		
Likelihood Ratio	.105	1	.746		
Fisher's Exact Test				.744	.487
Linear-by-Linear Association	.107	1	.744		
N of Valid Cases <sup>b</sup>	100				

a. 1 cells (25.0%) have expected count less than 5. The minimum expected count is 3.51.

b. Computed only for a 2x2 table

### Lampiran 7. Hasil SPSS Tingkat Kepatuhan Pasien Hipertensi

#### 1. Tingkat kepatuhan berdasarkan usia

##### Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
usia * kepatuhan	100	100.0%	0	.0%	100	100.0%

##### usia \* kepatuhan Crosstabulation

Count		kepatuhan		Total
		patuh	tidak patuh	
usia	kurang dari sama dengan 50 tahun	12	23	35
	lebih dari 50 tahun	31	34	65
Total		43	57	100

##### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	1.668 <sup>a</sup>	1	.196		
Continuity Correction <sup>b</sup>	1.166	1	.280		
Likelihood Ratio	1.688	1	.194		
Fisher's Exact Test				.212	.140
Linear-by-Linear Association	1.652	1	.199		
N of Valid Cases <sup>b</sup>	100				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 15.05.

b. Computed only for a 2x2 table

## 2. Tingkat kepatuhan berdasarkan jenis kelamin

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
JenisKelamin * kepatuhan	100	100.0%	0	.0%	100	100.0%

JenisKelamin \* kepatuhan Crosstabulation

Count		kepatuhan		
		patuh	tidak patuh	Total
JenisKelamin	pria	10	23	33
	wanita	33	34	67
Total		43	57	100

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	3.240 <sup>a</sup>	1	.072		
Continuity Correction <sup>b</sup>	2.513	1	.113		
Likelihood Ratio	3.311	1	.069		
Fisher's Exact Test				.088	.056
Linear-by-Linear Association	3.207	1	.073		
N of Valid Cases <sup>b</sup>	100				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 14.19.

b. Computed only for a 2x2 table

## 3. Tingkat kepatuhan berdasarkan jumlah antihipertensi

**Case Processing Summary**

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
JumlahAntihipertensi * kepatuhan	100	100.0%	0	.0%	100	100.0%

**JumlahAntihipertensi \* kepatuhan Crosstabulation**

Count		kepatuhan		Total
		patuh	tidak patuh	
		JumlahAntihipertensi	tunggal	
	kombinasi	20	31	51
Total		43	57	100

**Chi-Square Tests**

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.608 <sup>a</sup>	1	.435	.545	.282
Continuity Correction <sup>b</sup>	.334	1	.563		
Likelihood Ratio	.609	1	.435		
Fisher's Exact Test					
Linear-by-Linear Association	.602	1	.438		
N of Valid Cases <sup>b</sup>	100				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 21.07.

b. Computed only for a 2x2 table

## 4. Tingkat kepatuhan berdasarkan jumlah penyakit penyerta

**Case Processing Summary**

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
JumlahPenyakitPenyerta * kepatuhan	100	100.0%	0	.0%	100	100.0%

**JumlahPenyakitPenyerta \* kepatuhan Crosstabulation**

Count		kepatuhan		
		patuh	tidak patuh	Total
		JumlahPenyakitPenyerta	0	29
	1+	14	15	29
Total		43	57	100

**Chi-Square Tests**

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.464 <sup>a</sup>	1	.496		
Continuity Correction <sup>b</sup>	.210	1	.647		
Likelihood Ratio	.462	1	.497		
Fisher's Exact Test				.513	.322
Linear-by-Linear Association	.459	1	.498		
N of Valid Cases <sup>b</sup>	100				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 12.47.

b. Computed only for a 2x2 table

## 5. Tingkat kepatuhan berdasarkan pendidikan

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
pendidikan * kepatuhan	100	100.0%	0	.0%	100	100.0%

pendidikan \* kepatuhan Crosstabulation

Count		kepatuhan		
		patuh	tidak patuh	Total
		pendidikan kurang dari sama dengan SMP	26	36
pendidikan lebih dari SMP	17	21	38	
Total	43	57	100	

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.075 <sup>a</sup>	1	.784		
Continuity Correction <sup>b</sup>	.004	1	.947		
Likelihood Ratio	.075	1	.784		
Fisher's Exact Test				.837	.473
Linear-by-Linear Association	.075	1	.785		
N of Valid Cases <sup>b</sup>	100				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 16.34.

b. Computed only for a 2x2 table



## 6. Tingkat kepatuhan berdasarkan pekerjaan

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
pekerjaan * kepatuhan	100	100.0%	0	.0%	100	100.0%

pekerjaan \* kepatuhan Crosstabulation

Count		kepatuhan		Total
		patuh	tidak patuh	
pekerjaan	formal	8	19	27
	non formal	35	38	73
Total		43	57	100

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	2.698 <sup>a</sup>	1	.100		
Continuity Correction <sup>b</sup>	2.002	1	.157		
Likelihood Ratio	2.771	1	.096		
Fisher's Exact Test				.116	.077
Linear-by-Linear Association	2.671	1	.102		
N of Valid Cases <sup>b</sup>	100				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 11.61.

b. Computed only for a 2x2 table

**Lampiran 8.** Hasil SPSS Tingkat Kepatuhan berdasarkan Tingkat Pengetahuan**Case Processing Summary**

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
pengetahuan * kepatuhan	100	100.0%	0	.0%	100	100.0%

**pengetahuan \* kepatuhan Crosstabulation**

Count		kepatuhan		Total
		patuh	tidak patuh	
pengetahuan	tinggi	23	26	49
	rata-rata	12	19	31
	rendah	8	12	20
Total		43	57	100

**Chi-Square Tests**

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.616 <sup>a</sup>	2	.735
Likelihood Ratio	.617	2	.735
Linear-by-Linear Association	.427	1	.514
N of Valid Cases	100		

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 8.60.

**Lampiran 9.** Hasil regresi logistik hubungan kepercayaan dengan kepatuhan  
**Logistic Regression**

**Case Processing Summary**

Unweighted Cases <sup>a</sup>		N	Percent
Selected Cases	Included in Analysis	100	100.0
	Missing Cases	0	.0
	Total	100	100.0
Unselected Cases		0	.0
Total		100	100.0

a. If weight is in effect, see classification table for the total number of cases.

**Dependent Variable Encoding**

Original Value	Internal Value
patuh	0
tidak patuh	1

**Categorical Variables Codings**

		Frequency	Parameter coding
			(1)
GeneralHarmful	tinggi	87	1.000
	rendah	13	.000
SpecificNecessity	tinggi	76	1.000
	rendah	24	.000
SpescificConcerns	tinggi	60	1.000
	rendah	40	.000
GeneralOveruse	tinggi	55	1.000
	rendah	45	.000

## Block 0: Beginning Block

**Iteration History<sup>a,b,c</sup>**

Iteration		-2 Log likelihood	Coefficients
			Constant
Step 0	1	136.663	.280
	2	136.663	.282
	3	136.663	.282

- a. Constant is included in the model.
- b. Initial -2 Log Likelihood: 136.663
- c. Estimation terminated at iteration number 3 because parameter estimates changed by less than .001.

**Classification Table<sup>a,b</sup>**

Observed			Predicted		
			kepatuhan		Percentage Correct
			patuh	tidak patuh	
Step 0	kepatuhan	patuh	0	43	.0
		tidak patuh	0	57	100.0
	Overall Percentage				57.0

- a. Constant is included in the model.
- b. The cut value is .500

**Variables in the Equation**

	B	S.E.	Wald	df	Sig.	Exp(B)
Step 0 Constant	.282	.202	1.947	1	.163	1.326

## Variables not in the Equation

			Score	df	Sig.
Step 0	Variables	GeneralOveruse(1)	.910	1	.340
		SpecificNecessity(1)	.631	1	.427
		SpescificConcerns(1)	.823	1	.364
		GeneralHarmful(1)	.061	1	.805
Overall Statistics			2.470	4	.650

## Block 1: Method = Enter

Iteration History<sup>a,b,c,d</sup>

Iteration		-2 Log likelihood	Coefficients				
			Constant	GeneralOveruse(1)	SpecificNecessity(1)	SpescificConcerns(1)	GeneralHarmful(1)
Step 1	1	134.167	-.024	-.408	.396	-.278	.453
	2	134.160	-.025	-.428	.409	-.293	.477
	3	134.160	-.025	-.428	.409	-.293	.477

a. Method: Enter

b. Constant is included in the model.

c. Initial -2 Log Likelihood: 136.663

d. Estimation terminated at iteration number 3 because parameter estimates changed by less than .001.

## Omnibus Tests of Model Coefficients

		Chi-square	df	Sig.
Step 1	Step	2.503	4	.644
	Block	2.503	4	.644
	Model	2.503	4	.644

## Model Summary

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	134.160 <sup>a</sup>	.025	.033



## Variables in the Equation

	B	S.E.	Wald	df	Sig.	Exp(B)	95.0% C.I. for EXP(B)	
							Lower	Upper
							Step 1 <sup>a</sup> GeneralOveruse(1)	-.428
SpecificNecessity(1)	.409	.482	.720	1	.396	1.505	.585	3.873
SpescificConcerns(1)	-.293	.464	.397	1	.529	.746	.300	1.855
GeneralHarmful(1)	.477	.652	.536	1	.464	1.612	.449	5.783
Constant	-.025	.680	.001	1	.970	.975		

a. Variable(s) entered on step 1: GeneralOveruse, SpecificNecessity, SpescificConcerns, GeneralHarmful.

## Correlation Matrix

	Constant	GeneralOveruse(1)	SpecificNecessity(1)	SpescificConcerns(1)	GeneralHarmful(1)
Step 1 Constant	1.000	.028	-.544	-.186	-.664
GeneralOveruse(1)	.028	1.000	-.160	-.301	-.242
SpecificNecessity(1)	-.544	-.160	1.000	.108	.030
SpescificConcerns(1)	-.186	-.301	.108	1.000	-.212
GeneralHarmful(1)	-.664	-.242	.030	-.212	1.000

**Lampiran 10.** Hasil uji validitas dan reliabilitas kuesioner HFQ

**Reliability**

**Scale: ALL VARIABLES**

**Case Processing Summary**

		N	%
Cases	Valid	30	100.0
	Excluded <sup>a</sup>	0	.0
	Total	30	100.0

a. Listwise deletion based on all variables in the procedure.

**Reliability Statistics**

Cronbach's Alpha	Cronbach's Alpha Based on Standardized Items	N of Items
.867	.871	9

**Item Statistics**

	Mean	Std. Deviation	N
P1	.8333	.37905	30
P2	.9333	.25371	30
P3	.6667	.47946	30
P4	.9000	.30513	30
P5	.8333	.37905	30
P6	.8333	.37905	30
P7	.7667	.43018	30
P8	.7667	.43018	30
P9	.7333	.44978	30



**Inter-Item Correlation Matrix**

	P1	P2	P3	P4	P5	P6	P7	P8	P9
P1	1.000	.239	.443	.745	.520	.280	.388	.599	.337
P2	.239	1.000	.094	.356	.239	.598	.484	.169	.443
P3	.443	.094	1.000	.236	.632	.063	.279	.613	.373
P4	.745	.356	.236	1.000	.447	.447	.604	.342	.553
P5	.520	.239	.632	.447	1.000	.280	.388	.599	.337
P6	.280	.598	.063	.447	.280	1.000	.811	.176	.742
P7	.388	.484	.279	.604	.388	.811	1.000	.255	.915
P8	.599	.169	.613	.342	.599	.176	.255	1.000	.380
P9	.337	.443	.373	.553	.337	.742	.915	.380	1.000

**Item-Total Statistics**

	Scale Mean if Item Deleted	Scale Variance if Item Deleted	Corrected Item- Total Correlation	Squared Multiple Correlation	Cronbach's Alpha if Item Deleted
P1	6.4333	4.875	.625	.773	.850
P2	6.3333	5.471	.445	.376	.866
P3	6.6000	4.800	.492	.645	.866
P4	6.3667	5.068	.658	.757	.850
P5	6.4333	4.875	.625	.645	.850
P6	6.4333	4.944	.580	.759	.855
P7	6.5000	4.534	.734	.912	.839
P8	6.5000	4.810	.566	.681	.856
P9	6.5333	4.464	.735	.911	.839

**Scale Statistics**

Mean	Variance	Std. Deviation	N of Items
7.2667	6.064	2.46259	9

## Lampiran 11. Hasil uji validitas dan reliabilitas kuesioner BMQ

### Reliability

#### Scale: ALL VARIABLES

**Case Processing Summary**

		N	%
Cases	Valid	30	100.0
	Excluded <sup>a</sup>	0	.0
	Total	30	100.0

a. Listwise deletion based on all variables in the procedure.

**Reliability Statistics**

Cronbach's Alpha	Cronbach's Alpha Based on Standardized Items	N of Items
.865	.867	9

**Item Statistics**

	Mean	Std. Deviation	N
P1	2.8667	1.00801	30
P2	3.5333	1.50249	30
P3	3.9667	1.12903	30
P4	3.0667	.98027	30
P5	3.8333	1.11675	30
P6	3.1000	1.24152	30
P7	3.8000	1.12648	30
P8	4.2667	.78492	30
P9	4.0000	1.11417	30

Inter-Item Correlation Matrix

	P1	P2	P3	P4	P5	P6	P7	P8	P9
P1	1.000	.413	.269	.428	.531	.920	.401	.352	.399
P2	.413	1.000	.783	.490	.630	.414	.371	.548	.185
P3	.269	.783	1.000	.376	.488	.248	.211	.244	.137
P4	.428	.490	.376	1.000	.325	.448	.262	.335	.347
P5	.531	.630	.488	.325	1.000	.560	.466	.328	.610
P6	.920	.414	.248	.448	.560	1.000	.459	.396	.424
P7	.401	.371	.211	.262	.466	.459	1.000	.530	.659
P8	.352	.548	.244	.335	.328	.396	.530	1.000	.118
P9	.399	.185	.137	.347	.610	.424	.659	.118	1.000

Item-Total Statistics

	Scale Mean if Item Deleted	Scale Variance if Item Deleted	Corrected Item-Total Correlation	Squared Multiple Correlation	Cronbach's Alpha if Item Deleted
P1	29.5667	39.909	.673	.851	.845
P2	28.9000	35.059	.685	.821	.844
P3	28.4667	40.878	.509	.665	.859
P4	29.3667	41.757	.535	.455	.856
P5	28.6000	38.179	.730	.725	.838
P6	29.3333	37.540	.685	.867	.842
P7	28.6333	39.895	.585	.688	.852
P8	28.1667	43.592	.510	.574	.860
P9	28.4333	41.082	.502	.746	.860

Scale Statistics

Mean	Variance	Std. Deviation	N of Items
32.4333	49.495	7.03530	9