

DAFTAR PUSTAKA

- Arini, F. A. 2010. *Pengukuran Antropometri dan Hubungannya dengan "Golden Standard" Persen Lemak Tubuh*. Bioelectrical Impedance Analysis: Studi Validasi Pada Anak Sekolah Dasar. Depok. Universitas Indonesia.
- Araujo, D. M., Cabral, C. P., de Arruda, K. G. I., Falcao, S. T. A. P., Silvia, D. A. 2012. Body fat assessment by bioelectrical impedance and its correlation with anthropometric indicators. *Nutricion Hospitalaria*. 27: 1999-2005.
- Deurenberg, P. 2001. Original Communication: The Validity of Predicted Body Fat Percentage from Body Mass Index and from Impedance in Samples of Five European Populations. *European Journal of Clinical Nutrition*. 55: 973-979.
- Duncann, S. 2007. *Physical Activity and Obesity in Children: Measurement, Association Recommendation*. Auckland University of Technology New Zealand.
- Freedman, D., Berenson., G. S. 2009. Relation of circumferences and skinfold thicknesses to lipid and insulin concentrations in children and adolescents The Bogalusa Heart Study. *The American Journal of Clinical Nutrition*. 69: 308-317.
- Freedmant, S. D., Ogden, C. L., Blanck, H. M., Borrud, L. G., Dietz, W. H. 2013. The Abilities of Body Mass Index and Skinfold Thicknesses to Identify Children with Low or Elevated Levels of Dual Energy X-Ray Absoprtiometry-Determined Body Fatness. *Journal Peditary*. 163(1): 160-166.
- Gibson., Rosalin. S. 205. *Principal of Nutritional Assessment. Second Adition*. New York. USA. Oxford University Press, Inc.

- Guedes, D. P. 2012. Clinical procedures used for analysis of the body composition. *Brazilian Journal of Kinanthropometry and Human Performance*. 15(1): 113-129.
- Hills., Andrew., King, N., Byrne, N. 2007. Children, Obesity, and Exercise: Prevention, treatment and management of childhood and adolescent obesity. New York. Roulledge.
- Kelsey, H. C., Behnam, S., Claudia, S., Raylene, A. R., Walter, H., Rick, C., Deborah, A. M. 2016. Association of body mass index (BMI) and percentage body fat among BMI-defined non-obese middle-aged individuals: Insights from a population-based Canadian sample. *Canadian Journal of Public Health*. 107(6): 520- 525.
- Mastria, A. 2014. Hubungan Persentase Lemak Tubuh dengan Total Body Water Mahasiswa Kedokteran Universitas Diponegoro Semarang. *Jurnal Media Medika Muda*.
- Moreno, L. A., Warnberg, J. 2008. Body fat distribution reference standards in Spanish adolescents: The AVENA Study. *International Journal of Obesity*.31: 1798-1805.
- Navti, K. L., Atanga, M. B., Niba, L. L. 2017. Associations of out of school physical activity, sedentary lifestyle and socioeconomic status with weight status and adiposity of Cameroon children. *BMC Obesity*. 4(35).
- Nooyens, A. C. J., Koppes, L. L. J., Visscher, T. L. S., Twisk, J. W. R., Kemper, H. C. G., Schuit, A. J., van Mechelen, W., Seidell, J. C. 2007. Adolescent skinfold thickness is a better predictor of high body fatness in adults than is body mass index: the Amsterdam Growth and Health Longitudinal Study. *American Journal Clinical Nutrition*. 85: 1533-1539.
- Nurzakiah., Endang, A., Ratu, A. D. S. 2010. *Faktor Risiko Obesitas pada Orang Dewasa Urban dan Rural*. Artikel Penelitian FK UI. 29-34.

- Ojo, G., Adetola, O. 2017. The Relationship between Skinfold Thickness and Body Mass Index in Estimating Body Fat Percentage on Bowen University Students. *International Biology Biomed J.* 3: 3.
- Pongchaiyakul, C., Kosulwat, V., Rajroongwasinkul, N., Charoenkiatkul, S., Thesuthammarat, K., Laopaiboon, M., Nguyen, T. V., Rajatanavin, R. 2004. Prediction of Percentage Body Fat in Rural Thai Population Using Simple Anthropometric Measurements. *Obesity Research.* 13(4).
- Pradeep, S. C. 2014. Assessment of Body Fat Percentage by Different Methods: A Comparative Study. *European Journal of Sports and Exercise Science.* 3(3):1-6.
- Robinson, R., Mario, F. L., Jorge, E. C., Katherine, G., Emilio, G., Diana, P. C., Andres, B., Hector, R. T., Jacqueline, S. 2016. Triceps and Subscapular Skinfold Thickness Percentiles and Cut-Offs for Overweight and Obesity in a Population-Based Sample of Schoolchildren and Adolescents in Bogota, Colombia. *Nutrients.* 8(595).
- Sarria, A., Garcia-Llop, L. A., Moreno, L. A., Fleta, J., Morellon, M. P., Bueno, M. 1988. Skinfold thickness measurements are better predictors of body percentage than body mass index in males Spanish children and adolescents. *European Journal of Clinical Nutrition.* 52: 573-576.
- Wang, J. 1994. Asians Have Lower Body Mass Index (BMI) but Higher Percent Body Fat than Do Whites: Comparisons of Anthropometric Measurement. *American Journal of Clinical Nutrition.* 60: 23-28.